



Synthesis of some novel 3- (4-ethoxyphenyl) -5- (4-substituted) -4,5-dihydro-1H-pyrazole derivatives as potent antioxidant agents

Lakshminarayanan B^{*1,2}, Kannappan N², Subburaju T¹, Kalaichelvan V K²

¹Department of Pharmaceutical Chemistry, Ahalia School of Pharmacy, Palakkad-678557, Kerala, India

²Department of Pharmacy, Annamalai University, Chidambaram - 608002, Tamilnadu, India

Article History:

Received on: 19.09.2019

Revised on: 02.12.2019

Accepted on: 14.12.2019

Keywords:

Electron donating groups,
Electron withdrawing groups,
Ethoxy group,
In vitro antioxidant activity,
Pyrazoline,
Synthesis

ABSTRACT

Pyrazolines are the most useful heterocyclic moiety in Pharmaceutical and Chemical fields and as the most potential molecules for the design of new chemical entities. Nitrogen-containing heterocyclic compounds, pyrazolines and their derivatives showed a variety of pharmacological activities, including antioxidant properties. In the present study, eleven novel ethoxylated pyrazoline derivatives were synthesized by condensing chalcones with electron releasing ethoxy group at one end and different electron-donating, electron-withdrawing groups in another end with hydrazine hydrate and alcohol. The compounds synthesized were structural elucidated by their spectroscopic studies. All the compounds synthesized were evaluated for their *in vitro* antioxidant potential by 2,2'-diphenyl-1-picrylhydrazyl (DPPH) and hydrogen peroxide free radical scavenging assay methods. Some of these molecules possess moderate to good antioxidant activity when compared to standard ascorbic acid. The compound with methoxy group (EH2) exhibits potent antioxidant activity with IC₅₀ value of 9.02 and 9.44 μg/ml in DPPH and hydrogen peroxide assay methods respectively and the compound with hydroxy group (EH9) also showed potent antioxidant activity with IC₅₀ value of 12.41 and 14.56 μg/ml in DPPH and hydrogen peroxide free radical scavenging assay methods respectively when compared to standard. The compounds containing electron-donating substituents were found to be good antioxidants when compared to standard ascorbic acid.



*Corresponding Author

Name: Lakshminarayanan B

Phone: +91-9633769989

Email: blnrpharma@gmail.com

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v11i2.2035>

Production and Hosted by

Pharmascope.org

© 2020 | All rights reserved.

INTRODUCTION

Pyrazolines are well known five membered heterocyclic compounds with nitrogen and prepared

from α , β -unsaturated ketones with nucleophiles like NH_2NH_2 . H_2O with alcohol (Sakthnathan *et al.*, 2012; Sharath *et al.*, 2013). Pyrazoline with its derivatives are known to possess different biological and pharmacological activities like anti-inflammatory (Rathish *et al.*, 2009; Prabha *et al.*, 2019), antimicrobial (D'andrea *et al.*, 2005; Prabha *et al.*, 2019), antidepressant (Palaska *et al.*, 1996; Palaska, 2001), antitubercular (Kucukguzel and Rollas, 2002; Shaharyar *et al.*, 2006), antiviral (Kreutzberger and Kolter, 1986), antiamebic (Abid and Azam, 2005; Abid and Azam, 2006), anticonvulsant (Ozdemir *et al.*, 2007), analgesic (Amir *et al.*, 2008), anticancer (Taj *et al.*, 2011) and antioxidant potential (Martins *et al.*, 2009; Kumar *et al.*, 2013).

Generation of free radicals from the reactions of the imbalance between oxidative stress and antioxidant homeostatic phenomenon, which can lead to many pathological conditions in the body (Ashok and Tiwari, 2001). An elevated level of oxidative stress is one of the reasons for many diseases. Many biological damage due to the reactions of free radicals like H_2O_2 , hydroxyl radicals (OH), oxygen anions (O_2) free radicals, which leads to the process of chain reactions. Much reactive free radical plays a major role in the pathological condition of many diseases like heart disease, cancer, diabetes, strokes, Parkinson's diseases and allergies (Süzen, 2007).

The electron-donating CH_3 cluster substituted in the second position of 2-pyrazoline showed greater antioxidant property, whereas the electron-withdrawing Group (EWG) and high electron-donating groups were not increase free radicals inhibition (Kumar et al., 2013). Indole based pyrazole with an electron-donating group (EDG) has been proved increased antioxidant activity than EWG (Sharath et al., 2013). The presence of 2-quinolone with EWG in a pyrazoline ring showed increased antioxidant activity (Kumar et al., 2016). Nitrogen-containing heterocyclic compounds with the electron-withdrawing nature of halogen groups showed greater antioxidant property (Hossain et al., 2009).




Figure 1: Graphical abstract of the study

On the basis of these considerations, present research work was done to synthesize 1H-pyrazole derivatives with various EDG and EWG and evaluate them for antioxidant potential by DPPH & H_2O_2 assay methods. Graphical abstract of the present study was given in Figure 1.


MATERIALS AND METHODS

Chemicals and reagents

Analytical grade of reagent & chemicals were used in this study. All para-substituted aldehydes, hydrogen hydrate and other chemicals were procured from Sigma-Aldrich private limited.

Synthesis and characterization of ethoxylated pyrazolines

Eleven compounds of ethoxylated chalcones were prepared and evaluated their biological activities and reported (Lakshminarayanan et al., 2019). Continuation of the research, all the titled compounds (EH1 – EH11) were obtained by condensation of appropriate chalcones (E1 – E11) with 1.35 ml of hydrazine hydrate in the presence of 60 ml methanol and 15 ml acetic acid. The above mixture refluxed gently in a water bath for 6 – 8 hours. Then cool and acidified with conc. HCl. The product obtained was filtered and washed with water and recrystallized by ethanol. The open capillary method was used to determine the melting point and they were uncorrected. The purity was checked with TLC (hexane: ethyl acetate, 3:7) and melting point test and the route for the synthesis of titled molecules were given in Scheme 1. All prepared molecules were characterized by their spectral studies.



Scheme 1: Synthetic route for 1H-pyrazole derivatives

3-(4-ethoxyphenyl)-5-phenyl-4,5-dihydro-1H-pyrazole (EH1)

Brown: 54.3 %; m.p: 180-182°C. 1H NMR (400 MHz, $CDCl_3$) δ : 1.45-1.42 (t, 3H, $J=12.0$ Hz, CH_3-CH_2-O-), 3.93-3.91 (d, 2H, $J=8.0$ Hz, pyrazole H-4), 3.81-3.79 (t, 1H, $J=8.0$ Hz, pyrazole H-5), 4.09-4.07 (q, 2H, $J=8.0$ Hz, CH_3-CH_2-O-), 6.94-6.91 (d, 1H, $J=12.0$ Hz, H_8 & H_{10}), 7.27-7.25 (t, 1H, $J=8.0$ Hz, H_{15}), 7.32 (t, 1H, $J=12.0$ Hz, H_{14} & H_{16}), 7.47 (d, 1H, $J=12.0$ Hz, H_{13} & H_{17}), 7.92 (d, 1H, $J=12.0$ Hz, H_7 & H_{11}), 9.96 (s, 1H, NH). ^{13}C -NMR (100 MHz, $CDCl_3$) δ : 14.69 ($O-CH_2CH_3$), 42.46 (C_4), 59.75

(C₅), 63.72 (O-CH₂CH₃), 114.56 (C₈ & C₁₀), 125.50 (C₁₅), 127.12 (C₁₃ & C₁₇), 127.51 (C₆), 128.13 (C₇ & C₁₁), 128.79 (C₁₄ & C₁₆). ESI-MS (*m/z*): Calculated- 266.33, Observed- 266.32.

3-(4-ethoxyphenyl)-5-(4-methoxyphenyl)-4,5-dihydro-1H-pyrazole (EH2)

Pale brown; Yield: 59.2 %; m.p: 219-221°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.45-1.42 (t, 3H, *J*=12.0 Hz, CH₃-CH₂-O-), 3.86 (s, 3H, *J*=10.0 Hz, OCH₃), 3.78 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.72 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.08 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.94-6.92 (d, 1H, *J*=8.0 Hz, H₈ & H₁₀), 7.28 (d, 1H, *J*=12.0 Hz, H₁₄ & H₁₆), 7.41 (d, 1H, *J*=12.0 Hz, H₁₃ & H₁₇), 7.94 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.92 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.16 (O-CH₂CH₃), 30.11 (C₄), 64.24 (O-CH₂CH₃), 114.85 (C₈ & C₁₀), 127.32 (C₁₃ & C₁₇), 128.58 (C₆), 129.31 (C₇ & C₁₁), 114.58 (C₁₄ & C₁₆), 130.44 (C₁₂). ESI-MS (*m/z*): Calculated- 296.36, Observed- 296.35.

3-(4-ethoxyphenyl)-5-(4-methylphenyl)-4,5-dihydro-1H-pyrazole (EH3)

Pale brown; Yield: 32.1 %; m.p: 194-197°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.45-1.42 (t, 3H, *J*=12.0 Hz, CH₃-CH₂-O-), 2.41-2.40 (s, 3H, *J*=4.0 Hz, CH₃), 3.35 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.75 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.09-4.07 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.91-6.94 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.12 (d, 1H, *J*=12.0 Hz, H₁₄ & H₁₆), 7.27 (d, 1H, *J*=12.0 Hz, H₁₃ & H₁₇), 7.66-7.63 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.94 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.09 (O-CH₂CH₃), 21.42 (CH₃), 42.86 (C₄), 60.04 (C₅), 64.06 (O-CH₂CH₃), 114.54 (C₈ & C₁₀), 125.93 (C₁₃ & C₁₇), 127.35 (C₆), 128.53 (C₇ & C₁₁), 129.02 (C₁₄ & C₁₆), 141.33 (C₁₂), 161.12 (C₉). ESI-MS (*m/z*): Calculated- 280.36, Observed- 280.35.

4-[3-(4-ethoxyphenyl)-4,5-dihydro-1H-pyrazol-5-yl]-N,N-dimethylaniline (EH4)

Brown; Yield: 38.0 %; m.p: 233-235°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.46-1.43 (t, 3H, *J*=12.0 Hz, CH₃-CH₂-O-), 2.42 (s, 3H, *J*=4.0 Hz, CH₃), 3.92 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.97 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.09-4.07 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.72 (d, 1H, *J*=12.0 Hz, H₁₄ & H₁₆), 6.99-6.96 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.09 (d, 1H, *J*=12.0 Hz, H₁₃ & H₁₇), 7.86-7.83 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.96 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.28 (O-CH₂CH₃), 41.51 (CH₃), 42.84 (C₄), 52.04 (C₅), 64.16 (O-CH₂CH₃), 111.47 (C₁₄ & C₁₆), 114.44 (C₈ & C₁₀), 129.93 (C₁₃ & C₁₇), 128.13 (C₆), 128.93 (C₇ & C₁₁), 131.32 (C₁₂), 162.12 (C₉). ESI-MS (*m/z*): Calculated- 309.40, Observed- 309.39.

3-(4-ethoxyphenyl)-5-(4-nitrophenyl)-4,5-dihydro-1H-pyrazole (EH5)

Yellow: 41.7 %; m.p: 181-183°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.47-1.44 (t, 3H, *J*=12.0 Hz, CH₃-CH₂-O-), 2.53 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.15 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.12 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.97 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.27 (d, 1H, *J*=12.0 Hz, H₁₃ & H₁₇), 7.76 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 8.23 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 14.27 (O-CH₂CH₃), 63.36 (O-CH₂CH₃), 114.27 (C₈ & C₁₀), 122.33 (C₁₃ & C₁₇), 124.13 (C₁₄ & C₁₆), 127.55 (C₆), 128.16 (C₇ & C₁₁), 146.68 (C₁₅), 148.38 (C₁₂), 153.44 (C₆), 160.32 (C₉). ESI-MS (*m/z*): Calculated- 311.33, Observed- 311.32.

5-(4-chlorophenyl)-3-(4-ethoxyphenyl)-4,5-dihydro-1H-pyrazole (EH6)

Brown; Yield: 42.6 %; m.p: 208-210°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.47-1.44 (t, 3H, *J*=12.0 Hz, CH₃-CH₂-O-), 4.07-4.05 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 4.09 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 4.11 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 6.91-6.94 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.16-7.18 (d, 1H, *J*=8.0 Hz, H₁₄ & H₁₆), 7.27-7.29 (d, 1H, *J*=8.0 Hz, H₁₃ & H₁₇), 7.68-7.65 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.97 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 14.69 (O-CH₂CH₃), 42.33 (C₄), 59.21 (C₅), 63.63 (O-CH₂CH₃), 114.61 (C₈ & C₁₀), 127.08 (C₁₃ & C₁₇), 128.16 (C₆), 128.97 (C₇ & C₁₁), 133.27 (C₁₅), 140.40 (C₁₂), 153.69 (C₃), 160.77 (C₉). ESI-MS (*m/z*): Calculated- 300.78, Observed- 300.77.

3-(4-ethoxyphenyl)-5-(4-fluorophenyl)-4,5-dihydro-1H-pyrazole (EH7)

Brown; Yield: 49.4 %; m.p: 201-203°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.43-1.40 (t, 3H, *J*=12.0 Hz, CH₃-CH₂-O-), 3.9 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.94 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.01-3.99 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.95 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.15 (d, 1H, *J*=8.0 Hz, H₁₃ & H₁₇), 7.47 (d, 1H, *J*=8.0 Hz, H₁₄ & H₁₆), 7.70 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.93 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.06 (O-CH₂CH₃), 42.95 (C₄), 61.40 (C₅), 64.70 (O-CH₂CH₃), 115.25 (C₈ & C₁₀), 115.80 (C₁₄ & C₁₆), 129.05 (C₆), 130.36 (C₇ & C₁₁), 130.47 (C₁₃ & C₁₇), 131.15 (C₁₂), 163.37 (C₉). ESI-MS (*m/z*): Calculated- 284.32, Observed- 284.31.

5-(4-bromophenyl)-3-(4-ethoxyphenyl)-4,5-dihydro-1H-pyrazole (EH8)

Pale yellow; Yield: 41.1 %; m.p: 163-165°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.45-1.43 (t, 3H, *J*=8.0 Hz, CH₃-CH₂-O-), 3.69-3.66 (d, 2H, *J*=12.0 Hz, pyrazole H-4), 3.73-3.71 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.05-4.03 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.93-6.90 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.10-7.08 (d, 1H, *J*=8.0 Hz, H₁₃ & H₁₇), 7.40-7.42 (d, 1H, *J*=8.0 Hz, H₁₄ & H₁₆), 7.67-7.64 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.94

(s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.11 (O-CH₂CH₃), 42.76 (C₄), 55.55 (C₅), 64.30 (O-CH₂CH₃), 114.38 (C₈ & C₁₀), 121.92 (C₁₅), 127.85 (C₁₃ & C₁₇), 128.75 (C₆), 129.85 (C₇ & C₁₁), 131.16 (C₁₄ & C₁₆), 141.05 (C₁₂), 155.16 (C₃), 161.40 (C₉). ESI-MS (*m/z*): Calculated- 345.23, Observed- 345.22.

4-[3-(4-ethoxyphenyl)-4,5-dihydro-1H-pyrazol-5-yl]phenol (EH9)

Brown; Yield: 52.6 %; m.p: 182-183°C ¹H NMR (400 MHz, CDCl₃) δ: 1.45-1.43 (t, 3H, *J*=8.0 Hz, CH₃-CH₂-O-), 3.72-3.70 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.77-3.74 (t, 1H, *J*=12.0 Hz, pyrazole H-5), 4.09-4.07 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.93-6.91 (d, 1H, *J*=8.0 Hz, H₁₄ & H₁₆), 7.26 (d, 1H, *J*=12.0 Hz, H₁₃ & H₁₇), 7.78 (d, 1H, *J*=8.0 Hz, H₈ & H₁₀), 7.94-7.92 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 8.03 (s, 1H, OH), 9.96 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.07 (O-CH₂CH₃), 58.94 (C₅), 64.16 (O-CH₂CH₃), 114.51 (C₈ & C₁₀), 114.64 (C₁₄ & C₁₆), 128.87 (C₁₃ & C₁₇), 130.79 (C₆), 130.94 (C₇ & C₁₁), 131.03 (C₁₂). ESI-MS (*m/z*): Calculated- 282.33, Observed- 282.32.

3-(4-ethoxyphenyl)-5-(4-ethylphenyl)-4,5-dihydro-1H-pyrazole (EH10)

Pale yellow; Yield: 61.8 %; m.p: 165-167°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.26-1.24 (t, 3H, *J*= 8.0 Hz, CH₃-CH₂), 1.44-1.42 (t, 3H, *J*=8.0 Hz, CH₃-CH₂-O-), 2.73-2.71 (q, 3H, *J*= 8.0 Hz, CH₃-CH₂), 3.61-3.59 (d, 2H, *J*=8.0 Hz, pyrazole H-4), 3.72-3.69 (t, 1H, *J*=12.0 Hz, pyrazole H-5), 4.05-4.03 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 7.05-7.03 (d, 1H, *J*=8.0 Hz, H₁₄ & H₁₆), 7.23 (d, 1H, *J*=12.0 Hz, H₁₃ & H₁₇), 7.08 (d, 1H, *J*=8.0 Hz, H₈ & H₁₀), 7.93-7.91 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.93 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 14.91 (O-CH₂CH₃), 15.22 (-CH₂CH₃), 28.13 (-CH₂CH₃), 41.71 (C₄), 55.76 (C₅), 64.25 (O-CH₂CH₃), 114.71 (C₈ & C₁₀), 125.32 (C₁₃ & C₁₇), 128.21 (C₆), 128.62 (C₇ & C₁₁), 129.65 (C₁₄ & C₁₆), 140.43 (C₁₂), 152.12 (C₃), 161.55 (C₉). ESI-MS (*m/z*): Calculated- 294.39, Observed- 294.38.

3-(4-ethoxyphenyl)-5-[4-(trifluoromethyl)phenyl]-4,5-dihydro-1H-pyrazole (EH11)

Pale brown; Yield: 60.9 %; m.p: 189-191°C. ¹H NMR (400 MHz, CDCl₃) δ: 1.44-1.42 (t, 3H, *J*=8.0 Hz, CH₃-CH₂-O-), 3.68-3.65 (d, 2H, *J*=12.0 Hz, pyrazole H-4), 3.53-3.51 (t, 1H, *J*=8.0 Hz, pyrazole H-5), 4.06-4.04 (q, 2H, *J*=8.0 Hz, CH₃-CH₂-O-), 6.99-6.96 (d, 1H, *J*=12.0 Hz, H₈ & H₁₀), 7.18-7.16 (d, 1H, *J*=8.0 Hz, H₁₃ & H₁₇), 7.41-7.43 (d, 1H, *J*=8.0 Hz, H₁₄ & H₁₆), 7.89-7.86 (d, 1H, *J*=12.0 Hz, H₇ & H₁₁), 9.98 (s, 1H, NH). ¹³C-NMR (100 MHz, CDCl₃) δ: 15.76 (O-CH₂CH₃), 42.66 (C₄), 52.35 (C₅), 64.33 (O-CH₂CH₃), 114.90 (C₈ & C₁₀), 124.57 (C₁₄ & C₁₆), 128.05 (C₆), 128.42

(C₇ & C₁₁), 129.46 (C₁₃ & C₁₇), 145.16 (C₁₂). ESI-MS (*m/z*): Calculated- 334.33, Observed- 334.32.

In vitro antioxidant activity

DPPH free radical scavenging activity

All titled molecules were selected for their *in vitro* antioxidant potential by scavenging of DPPH free radical (Kaushik *et al.*, 2016). Standard ascorbic acid and synthesized compounds were prepared in different concentrations from 10 to 50 µg/ml from the stock solution (100 µg/ml) with distilled carbinol. In an individual test tube, one ml of each compound solution was taken; 4 ml of 0.004% methanol solution of DPPH was added and shaken vigorously. Keep the mixtures in a dark room for 30 minutes at room temperature and measure the absorbance of all the solution at 517 nm using a UV-VISIBLE spectrophotometer (Shimadzu UV-1800). Percentage radical scavenging activity was calculated with the formula [(Ab₀ - Ab₁)/Ab₀] x 100, where Ab₀ is the absorbance of blank, Ab₁ is the absorbance of synthesized compounds/standard. The calibration curve graph was plotted between percent inhibition and concentrations of the test/standard to get the amount of antioxidants need to decrease 50% from the starting concentration of DPPH free radicals. Half maximal inhibition values were calculated from the calibration curve. (Malladi *et al.*, 2014; Kaushik *et al.*, 2015).

Hydrogen peroxide free radical scavenging activity

Hydrogen peroxide free radical scavenging activities of titled compounds were done by the described method with a small modification (Babu *et al.*, 2001; Jayaprakasha *et al.*, 2004). Solutions of 20 mM hydrogen peroxide were prepared in phosphate buffer saline (PBS) and adjust the pH to 7.4. Standard ascorbic acid and synthesized compounds were prepared in different concentrations from 10 - 50 µg/ml with distilled carbinol. Each compound (1 ml) was taken in test tubes and adds 2 ml of H₂O₂ solution to all test tubes. Absorbance was measured at 230 nm after 10 minutes against a blank. The % scavenging activity and IC₅₀ values measured and intended by using the formula mentioned in the DPPH assay method.

RESULTS AND DISCUSSION

All the newly synthesized compounds were confirmed by spectroscopic studies.

In vitro antioxidant potency


Newly synthesized compounds (EH1 - EH11) are subjected to evaluate for antioxidant activities by

Table 1: Antioxidant activities of synthesized compounds (EH1 – EH11) against DPPH radical scavenging method

Compounds	% DPPH radical scavenging activity					IC ₅₀ (µg/ml)
	10 µg/ml	20 µg/ml	30 µg/ml	40 µg/ml	50 µg/ml	
EH1	19.56	19.99	21.47	22.90	23.56	31.84
EH2	50.03	56.72	58.34	61.09	63.56	9.02
EH3	41.21	44.51	48.54	51.35	52.09	25.75
EH4	45.62	48.76	51.82	54.45	55.68	29.02
EH5	40.31	42.68	44.28	45.26	46.74	130.73
EH6	51.25	54.33	58.52	63.00	65.77	36.33
EH7	29.98	32.32	33.65	35.23	36.12	46.56
EH8	29.08	32.17	34.69	36.11	38.49	95.29
EH9	47.17	53.83	55.52	58.12	60.77	12.41
EH10	21.34	22.35	31.12	32.45	34.01	25.67
EH11	19.25	21.6	22.44	24.54	25.97	115.38
Ascorbic acid	46.34	58.90	63.22	67.91	72.76	8.81

Table 2: Antioxidant activities of synthesized compounds (EH1 –EH11) against H₂O₂ radical scavenging method

Compounds	% H ₂ O ₂ radical scavenging activity					IC ₅₀ (µg/ml)
	10 µg/ml	20 µg/ml	30 µg/ml	40 µg/ml	50 µg/ml	
EH1	40.45	43.09	45.21	46.82	48.00	33.52
EH2	48.61	55.27	59.15	60.21	62.36	9.44
EH3	20.01	22.38	24.51	26.19	27.18	29.53
EH4	37.23	39.02	42.54	44.56	46.32	30.51
EH5	40.01	44.07	46.29	49.52	52.04	117.98
EH6	49.98	51.23	53.51	55.28	57.20	44.70
EH7	18.12	19.90	21.14	22.21	23.25	57.51
EH8	10.04	12.33	14.32	16.48	18.76	100.74
EH9	41.15	50.56	55.34	58.76	60.46	14.56
EH10	40.90	47.17	52.29	55.02	57.04	23.69
EH11	21.91	22.08	23.51	24.19	26.98	102.67
Ascorbic acid	42.45	48.35	51.16	52.89	54.29	7.65

**Figure 2: Comparative IC₅₀ values of the synthesized compounds in both assay methods**

DPPH and H₂O₂ free radicals scavenging methods. Absorbance of DPPH was decreased at 517 nm in all the compounds and standard was observed. The DPPH free radical scavenges activity and 50 % inhibitory concentrations (IC₅₀) for each compound were calculated and depicted in Table 1.

Hydrogen peroxide free radicals are toxic because it may give hydroxyl free radical to the cells even though they were not very reactive. So removal of H₂O₂ radicals is need for antioxidant activities in the cell. The ability of scavenging H₂O₂ free radicals and 50 % inhibitory concentrations (IC₅₀) for each compound were calculated and depicted in Table 2.

All the compounds showed moderate to good antioxidant potency compared with ascorbic acid. From the results, the compounds EH2 & EH9 with an IC₅₀ value of 9.02 and 12.41 in DPPH method and 9.44 & 14.56 in H₂O₂ assay method respectively. These compounds showed very good antioxidant potency among the series when compared to standard ascorbic acid (IC₅₀ value of 8.81 in DPPH and 7.65 in H₂O₂ method). It also found from Figure 2 that the synthesized compounds (EH1 – EH11) showed their IC₅₀ values in the DPPH method proportionate to the IC₅₀ values in the hydrogen peroxide assay method.

All the synthesized compounds contains electron-donating –O-CH₂-CH₃ group at one end and different electron-donating and electron-withdrawing groups at another end. Antioxidant potentials of all compounds were influenced by the substituent on an aromatic ring in each compound. The better antioxidant activities of EH2 and EH9 may be owing to the presence of electron giving nature of -OCH₃ and -OH group. On the other hand, the introduction of electron retreating substituents like -Cl, -F, -Br, -CF₃, -NO₂ in compounds EH6, EH7, EH8, EH11 and EH5 has led to the lower the activity when compared with standard. Antioxidant activity of all the compounds may be owed to the presence of electron giving –O-CH₂-CH₃ on one of an aromatic circle in all compounds.

CONCLUSIONS

In conclusion, a new class of nitrogen-containing heterocyclic compounds with the ethoxy group were prepared and screened for their antioxidant potency. Among the compounds, EH2 and EH9 showed better antioxidant activity by both assay methods. Compounds with methoxy and hydroxy groups possess good antioxidant activities when compared to standard.

ACKNOWLEDGEMENT

The authors express their sincere thanks to Sophisticated Analytical Instruments Facility (SAIF), NMR research centre, IISC, Bangalore, Karnataka for providing NMR spectra.

REFERENCES

- Abid, M., Azam, A. 2005. 1-Substituted Thiocarbamoyl-3-Phenyl-2-Pyrazolines: Synthesis and In Vitro Antiamoebic Activities. *European Journal of Medicinal Chemistry*, 40(9):935–942.
- Abid and Azam 2006. Synthesis, characterization and anti amoebic activity of 1-(thiazolo[4,5-b]quinoxaline-2-yl)-3-phenyl-2-pyrazoline derivatives. *Bioorg Med Chem Lett*, 16(10):2812–2816.
- Amir, M., Kumar, H., Khan, S. A. 2008. Synthesis and pharmacological evaluation of pyrazoline derivatives as new anti-inflammatory and analgesic agents. *Bioorganic & Medicinal Chemistry Letters*, 18(3):918–922.
- Ashok, K., Tiwari 2001. Imbalance in antioxidant defence and human diseases: Multiple approaches of natural antioxidants therapy. *Current Science*, 81(9):1179–1187.
- Babu, B., Shylesh, B., Padikkala, J. 2001. Antioxidant and hepatoprotective effect of *Acanthus ilicifolius*. *Fitoterapia*, 72(3):300–302.
- D'andrea, S., Zheng, Z. B., Denbleyker, K., Fung-Tom, J. C., Yang, H., Clark, J., Bronson, J. 2005. Synthesis and antibacterial activity of dihydro-1,2-oxazine and 2-pyrazoline oxazolidinones: novel analogs of linezolid. *Bioorganic & Medicinal Chemistry Letters*, 15(11):2834–2839.
- Hossain, M. M., Shaha, S. K., Aziz, F. 2009. Antioxidant potential study of some synthesized N-heterocycles. *Bangladesh Medical Research Council Bulletin*, 35(2):49–52.
- Jayaprakasha, G. K., Rao, L., Sakariah, K. K. 2004. Antioxidant activities of flavidin in different in vitro model systems. *Bioorganic & Medicinal Chemistry*, 12(19):5141–5146.
- Kaushik, N., Kumar, N., Kumar, A. 2015. Synthesis of substituted 5-phenyl-1-(5-phenyl)-isoxazol-3-yl)-1H-tetrazole as antioxidant agents. *Journal of Advanced Scientific Research*, 6(1):14–19.
- Kaushik, N., Kumar, N., Kumar, A. 2016. Synthesis, antioxidant and antidiabetic activity of 1-[(5-substituted phenyl)-4,5-dihydro-1H-pyrazol-3-yl]-5-phenyl-1H-tetrazole. *Indian Journal of Pharmaceutical Sciences*, 78(3):352–359.
- Kreutzberger and Kolter 1986. Antiviral agents. 27. The aminomethinylation of 5-oxo-2-pyrazoline-3-carboxylic acid derivatives. *Arch Pharm (Weinheim)*, 319(1):18–25.
- Kucukguzel, Rollas, S. 2002. Synthesis, characterization of novel coupling products and 4-arylhydrazono-2-pyrazoline-5-ones as potential antimycobacterial agents. *Il Farmaco*, 57(7):1253–1256.
- Kumar, A., Fernandes, J., Kumar, P. 2016. Synthesis and biological evaluation of some novel pyrazoline incorporated 2-quinolones. *Research Journal of Pharmacy and Technology*, 9(12).
- Kumar, A., Varadaraj, B. G., Singla, R. K. 2013. Syn-

- thesis and evaluation of the antioxidant activity of novel 3,5-disubstituted-2-pyrazolines. *Bulletin of Faculty of Pharmacy*, 51(2):167–173.
- Lakshminarayanan, B., Baek, S. C., Lee, J. P., Kannappan, N., Mangiatordi, G. F., Nicolotti, O., Mathew, B. 2019. Ethoxylated Head of Chalcones as a New Class of Multi-Targeted MAO Inhibitors. *ChemistrySelect*, 4(21). pages 6614–6619.
- Malladi, S. R., Anisetti, R., R. P. 2014. A facile synthesis, in vitro anti-inflammatory and antioxidant activity of novel benzimidazolylpyrano. *Indian J Pharm Sci*, 76(6):510–518. 2,3-d[[1,3]thiazolocarboxonitriles.
- Martins, D. M., Torres, B. G., Spohr, P. R., Machado, P., Bonacorso, H. G., Zanatta, N., Emanuelli 2009. Antioxidant Potential of New Pyrazoline Derivatives to Prevent Oxidative Damage. *Basic & Clinical Pharmacology & Toxicology*, 104(2):107–112.
- Ozdemir, Z., Kandilci, H. B., Gümüşel, B., Caliş, U., B. A. 2007. Synthesis and studies on an antidepressant and anticonvulsant activities of some 3-(2-furyl)-pyrazoline derivatives. *Eur J Med Chem*, 42(3):373–379.
- Palaska, E. 2001. Synthesis and antidepressant activities of some 3,5-diphenyl-2-pyrazolines. *European Journal of Medicinal Chemistry*, 36(6):539–543.
- Palaska, E., Erol, D., Demirdamar, R. 1996. Synthesis and antidepressant activities of some 1,3,5-triphenyl-2-pyrazolines. *European Journal of Medicinal Chemistry*, 31(1):80005–80010.
- Prabha, T., Jubie, S., Selvamani, P., Latha, S. 2019. Dual evaluation of some novel chalcone annulated pyrazolines as anti-inflammatory and antimicrobial agents via in-silico target study on cyclooxygenase-2. *International Journal of Research in Pharmaceutical Sciences*, 10(4):3159–3169. & Thangavel Sivakumar.
- Rathish, I. G., Javed, K., Ahmad, S., Bano, S., Alam, M. S., Pillai, K. K., Bagchi, V. 2009. Synthesis and anti-inflammatory activity of some new 1,3,5-trisubstituted pyrazolines bearing benzene sulfonamide. *Bioorganic & Medicinal Chemistry Letters*, 19(1):255–258.
- Sakthinathan, S. P., Vanangamudi, G., Thirunarayanan, G. 2012. Synthesis, spectral studies and antimicrobial activities of some 2-naphthyl pyrazoline derivatives. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 95:693–700.
- Shaharyar, M., Siddiqui, A. A., Ali, M. A., Sriram, D., Y. P. 2006. Synthesis and in vitro antimycobacterial activity of N1-nicotinoyl-3-(40-hydroxy-30-methyl phenyl)-5-[(sub)phenyl]-2-pyrazolines. *Bioorg Med Chem Lett*, 16(15):3947–3949.
- Sharath, V., Kumar, H. V., Naik, N. 2013. Synthesis of novel indole based scaffolds holding pyrazole ring as anti-inflammatory and antioxidant agents. *Journal of Pharmacy Research*, 6(7):785–790.
- Süzen, S. 2007. Antioxidant Activities of Synthetic Indole Derivatives and Possible Activity Mechanisms. *Bioactive Heterocycles V*, pages 145–178.
- Taj, T., Kamble, R. R., Gireesh, T. M., Hunnur, R. K., Margankop, S. B. 2011. One-pot synthesis of pyrazoline derivatised carbazoles as antitubercular, anticancer agents, their DNA cleavage and antioxidant activities. *European Journal of Medicinal Chemistry*, 46(9):4366–4373.