




# INTERNATIONAL JOURNAL OF RESEARCH IN PHARMACEUTICAL SCIENCES

Published by Pharmascope Publications

Journal Home Page: [www.pharmascope.org/ijrps](http://www.pharmascope.org/ijrps)

## *In vitro* anti-diabetic and anti-bacterial activity of unexplored species of *Atuna*

Sayyid Fasal Jassim VT\*<sup>1</sup> and Gopal V<sup>2</sup><sup>1</sup>Research Scholar, PRIST University, Thanjavur-613 403, Tamil Nadu, India<sup>2</sup>Department of Pharmacognosy, College of Pharmacy, MTPG & RIHS, A Govt. of Puducherry Institution, Puducherry-605 006, India

Article History:	ABSTRACT	
Received on: 21.08.2017 Revised on: 15.12.2017 Accepted on: 03.02.2018	<p>The present study highlights the Antidiabetic and Antibacterial activity of the various parts of wild plants <i>Atuna indica</i> (Bedd.) Kosterm and <i>Atuna travancorica</i> (Bedd.) Kosterm belonging to the family Chrysobalanaceae from Western Ghats, India. The leaves and flowers of <i>Atuna indica</i> and the leaves of <i>Atuna travancorica</i> were collected, shade dried, powdered and extracted in 50% aqueous ethanol by maceration process and concentrated to dryness with the use of rotary vacuum evaporator. The ethanol extracts of the test plant parts of the two species were subjected to the <i>In vitro</i> Antidiabetic and Antibacterial activity with the help of <math>\alpha</math>-amylase inhibition and microtiter plate assay methods respectively. The leaves of the two species possess Antidiabetic and Antibacterial activity; in particular leaves of <i>A.indica</i> and <i>A.travancorica</i> shows prominent Antidiabetic activity when compared with the other parts of the two species examined. In the case of Antibacterial activity the leaves of <i>A.travancorica</i> possess significant activity against <i>Pseudomonas aeruginosa</i> and <i>Bacillus subtilis</i>. The results suggested that the flowers of <i>A.indica</i> does not have the said activity. Based on the results obtained from the preliminary screening, it can be concluded that the leaves of <i>A.travancorica</i> has the maximum activity when compared with the other parts of the two species.</p>	
<b>Keywords:</b>		
<i>Atuna indica</i> , <i>Atuna travancorica</i> , $\alpha$ -amylase inhibition, Microtiter plate method		

\* Corresponding Author

Name: Sayyid Fasal Jassim VT

Phone: +91-9946024430

Email: sayyidjasim@gmail.com

ISSN: 0975-7538

DOI: <https://doi.org/10.26452/ijrps.v9i2.1510>

Production and Hosted by

Pharmascope.org

© 2018 Pharmascope Publications. All rights reserved.

### INTRODUCTION

The two species of *Atuna* distributed in the Southern Western Ghats are *Atuna indica* (Bedd.) Kosterm and *Atuna travancorica* (Bedd.) Kosterm. The specimens of *Atuna indica* were collected from Nadukani (Nilambur North Forest Division, Kerala State, India) and it was first collected by Beddomei


from Carcoorghats of Wayanad, and it was the re-discovery of this species, about 150 years later. Similarly, *Atuna travancorica* was described based on Hooker's collection, collected from Near the valara water falls, under Munnar Forest Division, Kerala State, India. The taxonomical as well as morphological characterization of the species was clearly done (Sasidharan and Sujanal, 2011).

*Atuna indica*, a 20m tall tree with smooth, thin, brown bark having elliptic-oblong or elliptic ovate leaves with 17-21cm long and 5.5-7.5cm wide with white flowers. Similarly, in the case of *Atuna travancorica* 25m tall tree with smooth thin grayish brown bark having alternate, lanceolate leaves with 7-16.5cm long and 1.7-4.5cm wide with pale lavender or white flowers. A unique feature of the family (Chrysobalanaceae) is seen such as they are prominently keeled (Ghillea *et al.*, 2003; Flowering Plants, 1989). In the case of *Atuna in-*


*dica*(Bedd.)Kosterm; Flowering and Fruiting: November to February; Habitat: West Coast tropical evergreen forests (Western Ghats in India) where as in the case of *Atuna travancorica*(Bedd.) Kosterm; Flowering and Fruiting: January to May; Habitat : West Coast tropical evergreen forests, usually riparian (Southern India, Travancore region) Both the species are belonging to the endangered category (Sudhakaret *al.*, 2007; Sasi-dharanN, 2002; Malin *et al.*, 2015; Jean *et al.*, 2003)

Diabetes is a serious metabolic disorder damaging many of the body systems such as the blood vessels and nerves and is the commonest endocrine disorder that affects approximately more than 100 million people worldwide. It is caused mainly due to the improper functioning of the pancreas with regard to the production of insulin leading to the abnormal blood glucose level.

Plants containing natural antioxidants such as flavonoids, tannins, C & E Vitamins stimulate the beta cells of pancreas thus reducing the blood glucose level. Also generally it is noticed that medicinal plants are cost effective devoid of side effects.(Kooti W *et al.*, 2016; Bhojar P *et al.*, 2012). Plants are a rich source of secondary metabolites such as flavonoids, alkaloids, tannins and terpenoids which have been found in vitro to have antimicrobial properties and their activity is probably due to their ability to complex with extracellular and soluble proteins and the ability to complex with bacterial cell walls.(Cowan MM, 1999; Palombo EA, 2011). In the species of *Atuna indica*, shows the presence of Umbelliferone, an active coumarin having a number of reported pharmacological activities found to possess therapeutically better antioxidant activity as reported (Asish *et al.*, 2013). In this context, the search to find the Antidiabetic and Antibacterial activities present in this wild species, *Atuna indica* and *Atuna travancorica* has been taken as the main objective of this study.



**Figure 1: Habit of *Atuna indica* (Bedd.) Kosterm**



**Figure 2: Habit of *Atuna travancorica* (Bedd.) Kosterm**

## MATERIALS AND METHODS

### Plant Materials

leaves and flowers of *Atuna indica* and leaves of *Atuna travancorica* were collected from Western Ghats of Kerala; *Atuna indica* collected from Nadukani (Nilambur North Forest Division, Kerala State, India) and *Atuna travancorica* collected from Near the valara waterfalls, under Munnar Forest Division, Kerala State, India.

### Extraction of Plant Parts

The collected parts viz., leaves and flowers of the two species were chopped into suitable smaller size and shade dried, powdered in a mixer and stored in suitable air tight containers. Weighed a quantity of 10 gram in three separate stoppered conical flasks with proper labeling on it. To that added 100 ml of 50% aqueous ethanol solution and kept aside with occasional shaking for a period of 24 Hours. After a period of 24 hours the supernatant solvent replaced with fresh 100 ml 50% aqueous ethanol solution and the mixture again kept for a period of 24 Hours.(Molnar *et al.*, 2017). After the completion of the prescribed time, both the supernatant liquids mixed together and concentrated for the complete removal of the solvent using rotary vacuum evaporator. The extracts obtained from leaves and flowers of each species properly labeled and transferred to air tight vials.

### Antidiabetic activity by $\alpha$ -Amylase Inhibition Assay

$\alpha$ -amylase inhibition activity was performed according to the modified method described by (Giancarlo *et al.*, 2006), Reagents Required: 0.5% (w/v) soluble starch solution, Colour reagent solution: 96mM 3.5 dinitro salicylic acid (20ml), 5.31M sodium potassium tartarate in 2M sodium hydroxide (8ml) and deionized water(12ml),  $\alpha$ - amylase solution (0.5unit/ml) 0.001g enzyme in 100ml of 20mM sodium phosphate buffer(pH-6.9).

Method: 25 $\mu$ L, 50 $\mu$ L, 100 $\mu$ L of extract and 1000 $\mu$ L of  $\alpha$ -amylase enzyme were mixed and incubated at 25 $^{\circ}$ C for 30 minutes. To all the test tubes added 1000 $\mu$ L of starch solution and the tube incubated at 25 $^{\circ}$ C for 3 minutes. Then, 1ml of the color reagent was added and the tubes were placed in water bath at 85 $^{\circ}$ C. After 15min, the reaction mixture was removed from the water bath and cooled thereafter; diluted with 9ml distilled water and absorbance value was determined at 540nm. The inhibition percentage of  $\alpha$ -amylase was assessed by the formula given below:

$$\text{Inhibition\%} = \frac{\text{AC540} - \text{AT540} \times 100}{\text{AC540}}$$

Where AC is Absorbance of control and AT is Absorbance of Test.

**Table 1: Antidiabetic activity of various extracts of *Atuna indica* (Bedd.) Kosterm and *Atuna travancorica* (Bedd.) Kosterm**

Name of the part	Sample No	Concentration (mg)	OD at 518nm	% of inhibition
<i>Leaves of Atuna indica</i>	1	0.5	0.328	18.78
		1	0.338	22.75
		5	0.341	<b>34.13</b>
<i>Flowers of Atuna indica</i>	2	0.5	0.308	20.90
		1	0.319	23.81
		5	0.324	28.57
<i>Leaves of Atuna travancorica</i>	3	0.5	0.299	26.46
		1	0.304	32.28
		5	0.313	<b>43.39</b>
<b>Control</b>	Nil	<b>0.378</b>	<b>Nil</b>	<b>Control</b>

**Table 2: Antibacterial activity of various extracts of *A. indica* and *A. travancorica***

Name of the Plant Part	Name of the Bacteria	% Inhibition					
		Control OD	Concentration (100mg/ml)				
			T1(10µL)	T2(15µL)	T3(20µL)	T4(25µL)	T5(50µL)
<i>Leaves of Atuna indica</i>	<i>E. coli</i>	1.288	18.40	21.51	23.52	30.12	31.75
	<i>Pseudomonas aeruginosa</i>	1.45	11.38	25.31	28.34	29.93	39.31
	<i>Bacillus subtilis</i>	1.547	28.64	30.25	33.68	37.10	38.20
	<i>Staphylococcus aureus</i>	1.259	8.02	11.91	17.95	23.19	29.07
<i>Flowers of Atuna indica</i>	<i>E. coli</i>	1.288	23.99	30.36	31.21	34.01	37.34
	<i>Pseudomonas aeruginosa</i>	1.45	5.86	9.52	10.90	13.52	15.45
	<i>Bacillus subtilis</i>	1.547	14.35	22.43	24.76	26.96	28.25
	<i>Staphylococcus aureus</i>	1.259	3.73	9.45	12.15	15.25	19.62
<i>Leaves of Atuna travancorica</i>	<i>E. coli</i>	1.288	26.48	33.39	40.61	46.20	50.16
	<i>Pseudomonas aeruginosa</i>	1.45	20.76	27.86	39.86	49.52	<b>55.24</b>
	<i>Bacillus subtilis</i>	1.547	24.95	29.09	33.81	38.66	<b>54.75</b>
	<i>Staphylococcus aureus</i>	1.259	17.79	19.70	23.59	30.02	37.97

**Antibacterial assay using Microtiter Plate Method**

Preparation of microbial culture: Using aseptic techniques, a single pure colony was transferred into a 10 ml of nutrient broth/potato dextrose broth, capped and placed in incubator overnight at 37°C. After incubation, using aseptic preparation, turbidity of suspensions was calculated and adjusted using McFarland standards as a reference. (Wiegand *et al.*, 2008)

Preparation of the microtiter plate plates: Microtitre plates were prepared under aseptic conditions. A sterile 96 well plate was labeled. A volume of 10µl, 50µl, and 100µl of test material was pipetted into the wells. 100µl of nutrient broth/potato dextrose broth was added to each well. Finally, 100µl of microbial suspension was added to each well. Control dilutions of test material were also kept. Plate was wrapped loosely with cling film to ensure that organism did not become dehydrated. Each plate had a set of controls: a column with all solutions except the test compound, and a column

with all solutions except the organism adding 100µl of nutrient broth/potato dextrose broth instead. The plates were incubated at 37°C for 24 hours and OD reading was taken (OD600) after sufficient incubation. Optical density (OD) was obtained from subtracting the control OD from the sample OD.

The % of inhibition was calculated from the following equation:

$$\% \text{ of inhibition} = \frac{\text{Control} - \text{Test}}{\text{Control}} \times 100$$

## RESULTS AND DISCUSSION

In the present study, each of the three samples weighed a quantity of amount 10gram in three separate and clearly labeled stoppered 250ml conical flask contains the powdered form of leaves and flowers of *Atuna indica* and leaves of *Atuna travancorica* in 100ml of 50% aqueous ethanol solution as solvent used for the extraction. From each sample, ~2gram extracts were collected.

**Antidiabetic activity:** In the case of Antidiabetic activity studies, *Atuna travancorica* leaves showed comparatively better activity with a percentage of alpha amylase inhibition of 43.39% at a concentration of 5mg whereas *Atuna indica* leaves showed an inhibition of 34.13% in the same concentration.

**Antibacterial activity:** In the case of Antibacterial activity studies, *Atuna travancorica* leaves showed comparatively better activity against both strains of bacteria *Pseudomonas aeruginosa* and *Bacillus subtilis* with a percentage inhibition of 55.24% and 54.75% respectively.

## CONCLUSION

The current study concluded that there is a remarkable Antidiabetic and Antibacterial activities in both the endangered tree species of *Atuna*. Based on the observation of the results, it can be appreciated that the Antidiabetic activity is prominent in the leaves of *Atuna indica* and *Atuna travancorica* and the Antibacterial activity is pronounced in the leaves of *Atuna travancorica*. Overall results is the leaves of *A. travancorica* shows prominent Antidiabetic and Antibacterial activities when compared with the other parts of the two species examined. From this studies, it is observed that there is an immense requirement for a detailed and authoritative phytochemical screening about the phytoconstituents present in the leaves of *Atuna indica* and *Atuna travancorica* by using modern analytical instruments to find out the mechanism based activity.

## REFERENCES

- Ahmed YA, Nasiruddin AK, Huda K. *The Islamic Guideline on Medicine*, Darussalam Publishers and Distributers, Portland, USA, 2010, p12-15.
- Andrews JM. Determination of minimum inhibitory concentrations. *J.Antimicrob.Chemther.*48 (Suppl.1) :5-16(2001).
- Ankur P, Makarand P, Radhika J, Evaluation of Anti-Diabetic Property of Extracts of Different Plant Parts of *Salacia chinensis* Linn, *J Biodivers Bio-pros Dev.*, 2014,1:1.
- Asish GR, Deepak M, George S, Indira B, Phytochemical Profiling & Antioxidant Activity of *Atunaindica* (Bedd.) Kosterm-An Unexplored Tree Species Reported from Western Ghats, India. *IJPPR*, 5(1), 2013, 27-30.
- Bhojar P, Burde VV, Baheti JR, Anti-Diabetic Potential of Herbal Medicines: A Review, *IJPRD*,2011; Vol4(1): March-2012, 67-80.
- Cowan MM, Plant Products as Antimicrobial Agents, *Clinical Microbiology Reviews*, 12(4), Oct. 1999, p. 564-582.
- Evanilson AF, Xavier HS, Randau KP, Chryso-balanaceae: traditional uses phytochemistry and pharmacology, *Brazilian Journal of Pharmacognosy*, 22(5), Sep./Oct. 2012, 1181-1186.
- Ghilleen T, Prance, Cyathia A et al., *Species Plantarum, Flora of the World, Part 10. Chryso-balanaceae*, Australian Biological Resources Study, Canberra, 2003, 66-75.
- Giancarlo, S., Rosa, M.L., Nadjafi, F., Francesco, M., 2006. Hypoglycaemic activity of two spices extracts: *Rhus coriaria* L. and *Bunium persicum* Boiss. *Nat. Prod. Res.*20, 882-886.
- Jassim SF, Gopal V. Conservation, Phytochemical Characterization and Pharmacological Evaluation of the Endangered Tree Species of *Atuna indica* and *Atuna travancorica* (Chryso-balanaceae), *International Journal of Phytopharmacology*, 8(3), 2017, 108-111.
- Jean PP, Priya D, Jean-Pierre P, Ramesh BR, Analysis of threatened endemic trees of the Western Ghats of India sheds new light on the Red Data Book of Indian Plants. *Biodiversity and Conservation*, 12, 2003, 2091-2106.
- Kooti1 W, Farokhipour M, Asadzadeh Z, Larky DA, Samani MA, The role of medicinal plants in the treatment of diabetes: a systematic review, *Electronic physician*, January 2016, Volume: 8, Issue: 1, Pages: 1832-1842.

- Malin R, Kirsty S, Emity B, Meirion J. Conserving the World's Most Threatened Trees, A Global Survey of exsitu Collections, 2015, 38.
- Maruthupandian A, Mohan VR, Kottaimuthu R, Ethnomedicinal plants used for the treatment of diabetes and jaundice by Palliyar tribals in Sirumalai hills, Western Ghats, Tamil Nadu, India, *Indian Journal of Natural Products and Resources*, Vol. 2(4), December 2011, pp. 493-497.
- Modak M, Priyanjali D, Jayant L, Ghaskadbi S, Thomas PAD, Indian Herbs and Herbal Drugs Used for the Treatment of Diabetes, *J. Clin. Biochem. Nutr.*, 40, 163-173, May 2007.
- Molnar M, Mendešević N, Šubarić D, Banjari I, Jokić S, Comparison of various Techniques for the extraction of Umbelliferone and harniarin in *Matricaria chamomilla* Processing fractions, *Chemistry Central Journal*(2017)11:78.
- Murugan T, Albino JW, Murugan M, Antimicrobial Activity and Phytochemical Constituents of Leaf Extracts of *Cassia auriculata*, *Indian J Pharm Sci.*, 2013 Jan-Feb; 75(1): 122-125.
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities, *Nature*, 403: 853-858.
- Nayar TS, Sibi M, Rasiya AB, Mohanan N, Rajkumar G, Flowering Plants of Kerala: Status and Statistics, *Rheedea*, Vol. 18(2), 95-106, 2008.
- Nono RN, Barboni L, Teponno RB, Quassinti L, Bramucci M et al., Antimicrobial, antioxidant, anti-inflammatory activities and phytoconstituents of extracts from the roots of *Dissotis thollonii* Cogn. (Melastomataceae), *South African Journal of Botany*, 93, (2014), 19-26.
- Otun1 KO, Olatunji GA, Ajiboye AT, Badeggi UM, Isolation and Characterization of the Chemical Constituents of the Stem Bark of *Parinari polyandra* Benth, *International Research Journal of Pure & Applied Chemistry*, 4(6), 2014, 710-717.
- Palombo EA, Traditional Medicinal Plant Extracts and Natural Products with Activity against Oral Bacteria: Potential Application in the Prevention and Treatment of Oral Diseases, *Evidence-Based Complementary and Alternative Medicine*, Volume 2011,1-15.
- Pascal JP, Ramesh BR, Franceschi DD, Wet evergreen forest types of the southern western ghats, India, *Tropical Ecology*, 45(2): 281-292, 2004.
- Remington. The Science and Practice of Pharmacy. 21<sup>st</sup>Edition, p87-90.
- Sasidharan N, Sujanalpal P. The genus *Atuna* (Chrysobalanaceae) in Southern Western Ghats, India. *Rheedea*, 21(1), 2011, 81-83.
- Sasidharan N. Floristic Studies in Parambikulam Wildlife Sanctuary, KFRRI Research Report No.246, ISSN 0970-8103, November 2002, 15-17.
- Satoskar RS, Nirmala NR, Bhandarkar SD. Pharmacology and Pharmacotherapeutics 2015; 24<sup>th</sup> Edition, p1-2.
- Sudhakar RC, Chiranjibi P, Reddy KN, Raju VS. Census of Endemic Flowering Plants of Kerala, India. *Journal of Plant Sciences*, 2(5), 2007, 489-503.
- Volga VR, Narayanan MKR, Kumar NA, Endemic Trees of Western Ghats-A Check List From Wayanad District, Kerala, India, *IJPAES*, Volume 3, Issue 2, April-June, 2013.
- Wadkar KA, Magdum CS, Patil SS, Naikwade NS, Anti-Diabetic Potential and Indian medicinal Plants, *Journal of Herbal Medicine and Toxicology*, 2 (1) 45-50 (2008).
- Wiegand I, Hilpert K, Hancock R. Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nature Protocols*,3(2), 2008,163-175.